



PCT






WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07D 401/12</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/57958</b> <b>(43) International Publication Date:</b> 23 December 1998 (23.12.98)
<b>(21) International Application Number:</b> PCT/US98/11403 <b>(22) International Filing Date:</b> 15 June 1998 (15.06.98) <b>(30) Priority Data:</b> 08/882,753      16 June 1997 (16.06.97)      US <b>(71) Applicant:</b> SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US). <b>(72) Inventors:</b> CHEN, Xing; 12 Bradford Lane, Plainsboro, NJ 08536 (US). POIRIER, Marc; 114 Sunshine Court, Parlin, NJ 08859 (US). WONG, Yee-Shing; 26 Sherbrooke Drive, Florham Park, NJ 07932 (US). WU, Guang-Zhong; 88 Vanderveer Drive, Basking Ridge, NJ 07920 (US). <b>(74) Agents:</b> MAGATTI, Anita, W. et al.; Schering-Plough Corporation, Patent Dept., K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		<b>(81) Designated States:</b> AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>

**(54) Title:** SYNTHESIS OF INTERMEDIATES USEFUL IN PREPARING BROMO-SUBSTITUTED TRICYCLIC COMPOUNDS**(57) Abstract**

The invention relates to a process for preparing a compound of formula (I), comprising: (a) reacting 2,5-dibromo-3-methylpyridine with an amine of the formula  $\text{NHR}^5\text{R}^6$  to obtain an amide; (b) reacting the amide with a compound of formula (3) in the presence of a strong base to obtain a compound of formula (4); (c) converting a compound of formula (4) to the corresponding cyano compound or aldehyde; (d) reacting the cyano compound or aldehyde with a piperidine derivative to obtain an aldehyde or alcohol of formula (7a) or (7b), respectively; (e) cyclizing a compound of formula (7a) or (7b); wherein  $\text{R}^1$ - $\text{R}^7$  are as defined in the specification.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5


10     **SYNTHESIS OF INTERMEDIATES USEFUL IN PREPARING**  
          **BROMO-SUBSTITUTED TRICYCLIC COMPOUNDS**

**BACKGROUND OF THE INVENTION**

15                 This invention provides an improved process for preparing  
intermediates useful in the preparation of bromo-substituted tricyclic  
compounds known as antihistamines and as inhibitors of farnesyl  
protein transferase (FPT). In particular, the compounds of this invention  
are useful in the preparation of antihistamines such as those disclosed  
20     in U.S. Patent 5,151,423, and of FPT inhibitors disclosed in International  
Application No. PCT/US96/19603, filed December 19, 1996.

**SUMMARY OF THE INVENTION**

25                 This invention provides a process for preparing a  
compound of the formula




wherein:

          R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently selected from the group  
consisting of hydrogen and halo, provided that at least one of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>  
30     and R<sup>4</sup> is hydrogen and at least one of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> is halo; and  
          the dotted line represents an optional double bond;


-2-

comprising:


(a) reacting a compound of formula 1



- (i) with an amine of the formula  $\text{NHR}^5\text{R}^6$ , wherein  $\text{R}^5$  is hydrogen and  $\text{R}^6$  is  $\text{C}_1\text{-C}_6$  alkyl, aryl or heteroaryl;  $\text{R}^5$  is  $\text{C}_1\text{-C}_6$  alkyl, aryl or heteroaryl and  $\text{R}^6$  is hydrogen;  $\text{R}^5$  and  $\text{R}^6$  are independently selected from the group consisting of  $\text{C}_1\text{-C}_6$  alkyl and aryl; or  $\text{R}^5$  and  $\text{R}^6$ , together with the nitrogen to which they are attached, form a ring comprising 4 to 6 carbon atoms or comprising 3 to 5 carbon atoms and one hetero moiety selected from the group consisting of  $-\text{O}-$  and  $-\text{NR}^9-$ , wherein  $\text{R}^9$  is H,  $\text{C}_1\text{-C}_6$  alkyl or phenyl; in the presence of a palladium catalyst and carbon monoxide to obtain an amide of formula 2:




- (ii) with an alcohol of the formula  $\text{R}^{10}\text{OH}$ , wherein  $\text{R}^{10}$  is  $\text{C}_1\text{-C}_6$  lower alkyl or  $\text{C}_3\text{-C}_6$  cycloalkyl, in the presence of a palladium catalyst and carbon monoxide to obtain the ester of formula 2A




followed by reacting the compound of 2A with an amine of formula  $\text{NHR}^5\text{R}^6$  to obtain the amide of formula 2;

- (b) reacting the amide of formula 2 with a compound of formula 3




wherein  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  are as defined above and  $\text{R}^7$  is Cl or Br, in the presence of a strong base to obtain a compound of formula 4

-3-




(c)(i) converting a compound of formula 4 to a cyano compound of formula 5a




; or

5 (c)(ii) converting a compound of formula 4 or a cyano compound of formula 5a to an aldehyde of formula 5b





(d) reacting compound 5a or 5b with a piperidine derivative of formula 6



10

wherein L is a leaving group selected from the group consisting of Cl and Br, to obtain a ketone of formula 7a or an alcohol of formula 7b, respectively:



15

(e)(i) cyclizing a compound of formula 7a to obtain a compound of formula I wherein the dotted line represents a double bond; or

-4-

(e)(ii) cyclizing a compound of formula 7b to obtain a compound of formula I wherein the dotted line represents a single bond.

Preferred compounds of formula I are those wherein R<sup>2</sup> is halo. Also preferred are compounds wherein R<sup>1</sup> and R<sup>3</sup> are each hydrogen. Another group of preferred compounds is that wherein R<sup>1</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and R<sup>2</sup> is halo. Still another group of preferred compounds is that wherein R<sup>1</sup> and R<sup>3</sup> are each hydrogen and R<sup>2</sup> is halo. Yet another group of preferred compounds is that wherein R<sup>1</sup> and R<sup>3</sup> are each hydrogen and R<sup>2</sup> and R<sup>4</sup> are independently selected from the group consisting of halo. Halo is preferably Cl or Br.

#### **DETAILED DESCRIPTION**

As used herein, the term "lower alkyl" means straight or branched alkyl chains of 1 to 6 carbon atoms.

"Halo" refers to fluorine, chlorine, bromine or iodine radicals.

"Aryl" means phenyl, substituted phenyl wherein the substituents are 1 to 3 substituents independently selected from the group consisting of C<sub>1</sub> to C<sub>6</sub> alkyl and C<sub>1</sub> to C<sub>6</sub> alkoxy, benzyloxy or naphthyl.

"Heteroaryl" means a 5- or 6-membered aromatic ring comprising one or two nitrogen atoms, e.g., pyridyl, pyrimidyl, imidazolyl or pyrrolyl.

When R<sup>5</sup> and R<sup>6</sup>, together with the nitrogen to which they are attached, form a ring comprising 4 to 6 carbon atoms, the rings so produced are exemplified by pyrrolidinyl, piperidinyl and perhydroazepine. When R<sup>5</sup> and R<sup>6</sup>, together with the nitrogen to which they are attached, form a ring comprising 4 to 5 carbon atoms and a heteroatom, the rings so produced are exemplified by piperazinyl, N-methyl-piperazinyl, N-phenyl-piperazinyl and morpholinyl.

The compounds prepared by the process disclosed above are useful as intermediates in the procedures described in PCT/US96/19603 and U.S. 5,151,423 to obtain the desired compounds wherein the piperidinyl ring is N-substituted. By using the 3-bromo-substituted intermediates prepared by the process of this invention, the

-5-

desired tricyclic antihistamines and FPT inhibitors described above can be made by an eleven-step process rather than the fifteen-step process disclosed in the art.

Compounds of formula I can be converted to other compounds of formula I by methods known in the art, i.e., compounds wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup> is hydrogen can be converted to the corresponding compounds wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup> is halogen. Such procedures are shown in PCT/US96/19603, wherein, for example, a compound wherein R<sup>2</sup> is Cl, R<sup>1</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and the piperidinyl nitrogen is protected by a -COOCH<sub>2</sub>CH<sub>3</sub> group is reacted with KNO<sub>3</sub>, the resulting nitro-substituted compound is reduced to the amine, the resulting compound is reacted with Br<sub>2</sub> and the amino group is removed to obtain a compound wherein R<sup>2</sup> is Cl, R<sup>4</sup> is Br and R<sup>1</sup> and R<sup>3</sup> are hydrogen.

In step (a), the di-bromo-substituted pyridine of formula 1 is reacted with the amine NHR<sup>5</sup>R<sup>6</sup> in the presence of a palladium catalyst, carbon monoxide (CO) and a base. As defined above, the amines of formula NHR<sup>5</sup>R<sup>6</sup> are exemplified by t-butylamine, aniline, N-methylaniline, pyrrolidine, piperidine, perhydroazepine, piperazine, N-methylpiperazine, N-phenylpiperazine and morpholine. Preferred amines are pyrrolidine and t-butylamine, with t-butylamine being most preferred.

Palladium catalysts are exemplified by Pd(OAc)<sub>2</sub>/P(R<sup>11</sup>)<sub>3</sub> at ratios of 1:1 or 1:2; (PPh<sub>3</sub>)<sub>2</sub>PdCl<sub>2</sub> at a range of 0.5 to 40 mol%, preferably 1 to 10 mol%, and most preferably 1 to 5 mol%; Pd(PPh<sub>3</sub>)<sub>4</sub>; (R<sup>11</sup>)<sub>3</sub>P/Pd<sub>2</sub>(dba)<sub>3</sub>; Pd(OAc)<sub>2</sub>/2,2'-bipyridine at ratios of 1:1 to 1:2, preferably 1 to 10 mol%; and Pd/C, wherein Ac is acetyl, R<sup>11</sup> is C<sub>1</sub> to C<sub>6</sub> alkyl or aryl, Ph is phenyl, and dba is dibenzylidene acetone. Preferred catalysts are Pd(OAc)<sub>2</sub>/P(R<sup>11</sup>)<sub>3</sub> and (PPh<sub>3</sub>)<sub>2</sub>PdCl<sub>2</sub>.

The amount of amine (NHR<sup>5</sup>R<sup>6</sup>) reacted ranges from 1 to 4 equivalents, and is preferably 1 to 1.5 equivalents. Suitable bases include, but are not limited to, C<sub>1</sub> to C<sub>6</sub> alkyl amines such as triethylamine (Et<sub>3</sub>N), t-butylamine and 1,8-diazobicyclo[5.4.0]undec-7-ene (DBU), and inorganic bases such as K<sub>2</sub>CO<sub>3</sub>, Na<sub>2</sub>CO<sub>3</sub>, Na<sub>2</sub>HPO<sub>4</sub> and NaOH. Preferred bases are K<sub>2</sub>CO<sub>3</sub> and Et<sub>3</sub>N, with Et<sub>3</sub>N being most preferred.

-6-

Suitable solvents are tetrahydrofuran (THF), dimethylformamide (DMF), acetonitrile ( $\text{CH}_3\text{CN}$ ) and toluene or a combination thereof.  $\text{CH}_3\text{CN}$  is preferred for reaction with an amine and a combination of  $\text{CH}_3\text{CN}$  and toluene is preferred for reaction with an alcohol. The temperature range for the reaction is  $35^\circ\text{C}$  to  $100^\circ\text{C}$ , preferably about  $55^\circ\text{C}$  for reaction with the amine and preferably about  $80^\circ\text{C}$  for reaction with an alcohol. The reaction is carried out at a pressure of 5 psi to 500 psi, preferably 40 to 200 psi, and most preferably at 50 to 150 psi. The time for reaction ranges from 2 hours to 4 days, preferably 4 hours to 2 days, and most preferably 16 to 48 hours.

Conversion of the ester of formula 2A to the amide of formula 2 is accomplished by methods well known in the art, for example by reacting the ester directly with the amine or by using the conditions described by Basha et al in Tetrahedron Letters, (1977), p. 4171.

In step (b), the amide formed in step (a) is reacted with the halomethyl-substituted compound of formula 3 in a solvent such as THF, t-butyl methyl ether (t-BuOMe), diethyl ether ( $\text{Et}_2\text{O}$ ), diglyme or a mixture thereof, preferably a mixture of THF and t-butyl methyl ether, in the presence of a strong base such as lithium diisopropylamide (LDA), lithium hexamethyldisilylamide or sodium amide, preferably LDA. The concentration of the base ranges from 2.0 to 4.0 equivalents, preferably 2.0 to 2.2 equivalents. The compound of formula 3 is reacted in a concentration range of 1.0 to 1.5 equivalents, preferably 1.1 equivalents. The reaction is carried out in a temperature range of  $-78^\circ\text{C}$  to  $-20^\circ\text{C}$ , preferably  $-50^\circ\text{C}$  to  $-30^\circ\text{C}$ .

In step (c)(i), the product of step (b) is converted to the corresponding cyano compound of formula 5a by reacting with  $\text{POCl}_3$  or  $\text{SOCl}_2$  in a solvent such as  $\text{CH}_2\text{Cl}_2$  or without a solvent, preferably without a solvent. The reaction is carried out in a temperature range of  $50^\circ\text{C}$  to reflux, preferably at reflux.

Alternatively, in step (c)(ii), the product of step (b) or step (c)(i) is converted to the corresponding aldehyde of formula 5b by reacting with DIBALH or  $\text{LiAlH}_4$  and its derivatives, preferably DIBALH, in a solvent such as toluene, THF or t-BuOMe, preferably toluene. The



-7-

reaction is carried out in a temperature range of  $-78^{\circ}\text{C}$  to  $-30^{\circ}\text{C}$ , preferably  $-78^{\circ}\text{C}$  to  $-50^{\circ}\text{C}$ .

In step (d), the product of step 5a or 5b is reacted with a piperidine derivative of formula 6 as defined above to obtain a ketone or alcohol, respectively. The reaction is carried out in a solvent such as THF, toluene or t-BuOMe, preferably THF. The concentration of the piperidine derivative ranges from 1.0 to 2.0 equivalents, preferably 1.1 to 1.2 equivalents. The reaction is carried out in a temperature range from  $-20^{\circ}\text{C}$  to  $50^{\circ}\text{C}$ , preferably  $35^{\circ}\text{C}$  to  $45^{\circ}\text{C}$ , for the product of step 5a, and in a range from  $-78^{\circ}\text{C}$  to  $0^{\circ}\text{C}$ , preferably  $-78^{\circ}\text{C}$  to  $-60^{\circ}\text{C}$ , for the product of step 5b.

In step (e)(i), the ketone of formula 7a is cyclized to a compound of formula I wherein the dotted line represents a double bond by treatment with a strong acid such as  $\text{CF}_3\text{SO}_3\text{H}$ ,  $\text{CH}_3\text{SO}_3\text{H}$  or  $\text{BF}_3\cdot\text{HF}$ , preferably  $\text{CF}_3\text{SO}_3\text{H}$ , in a temperature range of  $50^{\circ}\text{C}$  to  $120^{\circ}\text{C}$ , preferably  $90^{\circ}\text{C}$  to  $95^{\circ}\text{C}$ .

In step (e)(ii), the alcohol of formula 7b is cyclized to a compound of formula I wherein the dotted line represents a single bond by treatment with an acid such as  $\text{H}_2\text{SO}_4$ , polyphosphoric acid or  $\text{CH}_3\text{SO}_3\text{H}$ , preferably polyphosphoric acid, in a temperature range of  $100^{\circ}\text{C}$  to  $200^{\circ}\text{C}$ , preferably  $160^{\circ}\text{C}$  to  $180^{\circ}\text{C}$ .

Starting materials of formula 1, 3, 6 and  $\text{NHR}^5\text{R}^6$  are known in the art or can readily be prepared by one skilled in the art.


Following are specific examples of the procedures in the various steps of the process of this invention for preparing compounds of formula I, although those skilled in the art will appreciate that similar procedures within the scope of the process of this invention can be used to prepare other compounds of formula I.

30


35

-8-

## EXAMPLE 1




Step (a):



5 To an autoclave were added 16 g (60.6 mmole) of 2,5-dibromo-3-methylpyridine, 4.5 g (6.4 mmole) of  $(\text{Ph}_3\text{P})_2\text{PdCl}_2$ , 150 ml of toluene, 150 ml of  $\text{CH}_3\text{CN}$ , and 17 ml (160 mmole) of *t*-butylamine. The autoclave was sealed, evacuated, purged with nitrogen and charged with carbon monoxide to 120 psi. The reaction mixture was heated to 60°C for two days with periodical refilling, as necessary, and then cooled to r.t. The contents of the autoclave was vented under vacuum, flushed with nitrogen and transferred to a flask with the aid of water and EtOAc. The mixture was concentrated and filtered through a pad of celite. The filtrate was extracted with EtOAc, the combined extract was washed with brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was separated by column chromatography to obtain 11 g of the product as an oil, 67 % yield.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 8.40 (d,  $J=2.1$ , 1H), 7.90 (br, 1H), 7.73 (d,  $J=2.1$ , 1H), 2.73 (s, 3H), 1.48 (s, 9H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ): 164.53, 146.58, 146.08, 142.92, 136.93, 122.28, 50.86, 28.71, 20.57.

Step (b):



To a solution of 10.56 ml (80.7 mmole) of  $i\text{-Pr}_2\text{NH}$  in 90 ml THF at 0°C was added 31.20 ml (77.9 mmole) of 2.5 M *n*-BuLi in hexanes and


-9-

the solution was stirred for 30 min. To the LDA solution was added dropwise a solution of 9.6 g (35.4 mmole) of the product of Step (a) in 45 ml THF at -78°C. The resulting purple solution was stirred at -78°C for 30 min., at -42°C for 15 min., and then recooled to -78°C. To this

5 solution was added dropwise a solution of 8.0 g (50 mmole) of 3-chlorobenzyl chloride in 50 ml THF. The reaction was warmed to room temperature over 1 hour. Saturated NH<sub>4</sub>Cl solution (50 ml) and ice-water (50 ml) were added to the reaction and the mixture was evaporated to half of the volume under vacuum. Extraction with EtOAc

10 (100 ml x 2) and evaporation of the solvent gave 16 g of the desired product, which was used directly in next step. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 8.40 (d, J=2.1, 1H), 7.78 (b, 1H), 7.55 (d, J=2.1, 1H), 7.16 (m, 3H), 7.06 (m, 1H), 3.38 (m, 2H), 2.90 (m, 2H), 1.45 (s, 9H).

Step (c)




The amide (16 g) of Step (b) was dissolved in POCl<sub>3</sub> (100ml) and the solution was refluxed for 2.5 hr and then concentrated to a third of its volume under vacuum, poured into 200 g ice and adjusted to pH 8 with 50% NaOH at 25°C. The resultant mixture was stirred for 2 hr at 25°C

20 and the pH maintained at 8 with NaOH. Extraction with EtOAc (100ml x 2) and evaporation gave a solid residue which was washed with hexane. After drying, 10 g of product was obtained; the yield was 88% in two steps. <sup>1</sup>NMR (CDCl<sub>3</sub>): 8.62 (d, J=2.0, 1H), 7.71 (d, J=2.0, 1H), 7.23 (m, 2H), 7.16 (s, 1H), 7.04 (m, 1H), 3.11 (m, 2H), 2.95 (m, 2H).

25 <sup>13</sup>C NMR (CDCl<sub>3</sub>): 150.21, 142.66, 141.10, 139.97, 134.42, 131.86, 129.96, 128.54, 126.95, 126.63, 124.60, 115.63, 35.94, 34.26.


-10-

Step (d)



To a solution of the product of Step (c) (2 g, 6.25 mmole) in THF (20 ml) at 40-45°C was added dropwise N-methyl-piperidyl magnesium chloride (8 ml, 0.94 M, 1.2 eq) and the reaction mixture was stirred for 30 min. The reaction mixture was adjusted to pH 2 with 2N HCl and was stirred for 1 h. The pH was adjusted to 10 with 28% NH<sub>4</sub>OH and the mixture was extracted with EtOAc (100ml x 2). The organic layer was separated and concentrated to give a residue, which was passed through silica gel as a CH<sub>2</sub>Cl<sub>2</sub> solution. The solvent was removed to obtain an oil (2.3 g). <sup>1</sup>H NMR (CDCl<sub>3</sub>): 8.54 (d, J=2.1, 1H), 7.62 (d, J=2.1, 1H), 7.08 (m, 3H), 7.30 (dt, J=7.0, 1.5, 1H), 3.62 (m, 1H), 3.08 (m, 2H), 2.86 (m, 4H), 2.28 (s, 3H), 2.06 (m, 2H), 1.82 (m, 2H), 1.75 (m, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>): 205.12, 150.09, 147.62, 142.81, 141.65, 139.37, 134.14, 129.69, 128.59, 126.75, 126.41, 123.17, 55.19, 46.35, 43.73, 36.97, 34.63, 27.93.

Step (e):




Stir a solution of ~2 g (4.6 mmole) of the product of Step (d) in 4.6 ml (50 mmole) of CF<sub>3</sub>SO<sub>3</sub>H at 90°C for 18 hr under nitrogen. Pour the cooled reaction into ice water and adjust to pH 10 with 29% NH<sub>4</sub>OH. Extract the product with CH<sub>2</sub>Cl<sub>2</sub> (2X) to obtain 2 g residue. Purify by chromatography on a silica gel column, eluting with CH<sub>2</sub>Cl<sub>2</sub>:CH<sub>3</sub>OH: NH<sub>4</sub>OH (28%) (100:3:0.1). The yield is 68% based on consumption of the starting ketone. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 8.44 (d, J=2.2, 1H), 7.56 (d, J=2.2,

-11-


1H), 7.12 (m, 3H), 3.36 (m, 2H), 2.70 (m, 4H), 2.50 (m, 1H), 2.35 (m, 3H), 2.25 (s, 3H), 2.05 (m, 2H).  $^{13}\text{C}$  ( $\text{CDCl}_3$ ): 155.79, 147.36, 139.57, 139.12, 137.45, 135.18, 132.78, 131.62, 130.64, 128.77, 126.12, 118.53, 56.74, 45.93, 31.33, 31.30, 30.99, 30.73.

5

## EXAMPLE 2




Step1:



To a solution of the starting amide (8 g, 20.3 mmole) in 80 ml of  
10 toluene at  $-70^\circ\text{C}$  was added 22 ml (22 mmole in toluene) of DIBALH  
dropwise over 10 min. The reaction was monitored by TLC; after  
completion, the reaction mixture was transferred at  $-60^\circ\text{C}$  to a quenching  
solution prepared with 150 ml of water and 11 g of malic acid, with the  
pH being adjusted to 14 with 50% NaOH. The resultant mixture was  
15 stirred for 15 min and the toluene layer was separated. The aqueous  
layer was extracted with toluene (100mL), the toluene layers were  
combined, dried over  $\text{MgSO}_4$  and filtered. Concentration gave 5.7 g  
product, 87% yield.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 10.10 (s, 1H), 8.71 (d,  $J=2.0$ , 1H),  
7.67 (d,  $J=2.0$ , 1H), 7.10-7.20 (m, 4H), 3.25 (m, 2H), 2.85 (m, 2H).


20 Step 2:



-12-

To a mixture of the aldehyde of Step 1 (0.32 g, 0.60mmol) in THF (20 ml) at -78°C was added dropwise the Grignard reagent (0.9 M, 0.7 ml, 0.63 mmol). After 30 min, aqueous NH<sub>4</sub>Cl (~2 ml) was added and the mixture was warmed to room temperature. Water (50ml) was added and the mixture was extracted with EtOAc (50ml x 2). After concentrating the combined organic phase, the residue was separated by preparative TLC to give 75 mg of product. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 8.49 (d, J=2.1, 1H), 7.66 (d, J=2.1, 1H), 7.51 (d, J=2.3, 1H), 7.05 (d, J=2.3, 1H), 4.69 (d, J=4.50, 1H), 2.70-3.10 (m, 6H), 2.19 (s, 3H), 1.80 (m, 3H), 1.55 (m, 4H), 1.35 (m, 1H).

Step 3:




A mixture of 0.5 g of the alcohol from Step 2 with 5 g of polyphosphoric acid was heated to 170°C for 2 hr. After cooling to r.t., the reaction was adjusted to pH 12 with aqueous NaOH and extracted with EtOAc. The organic layer was combined and dried over MgSO<sub>4</sub> and concentrated to give the product. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 8.35 (d, J=2.0, 1H), 7.50 (d, 1H), 7.06 (m, 3H), 3.85 (d, J=6.3, 1H), 3.35 (m, 2H), 2.80 (m, 4H), 2.20 (s, 3H), 2.05 (m, 1H), 1.75 (m, 2H), 1.20-1.50 (m, 4H).

20

-13-

We Claim:

1. A process for preparing a compound of the formula I




5

wherein:


R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of hydrogen and halo, provided that at least one of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> is hydrogen and at least one of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> is halo; and

- 10 the dotted line represents an optional double bond; comprising:

- (a) reacting a compound of formula 1




- 15 (i) with an amine of the formula NHR<sup>5</sup>R<sup>6</sup>, wherein R<sup>5</sup> is hydrogen and R<sup>6</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, aryl or heteroaryl; R<sup>5</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, aryl or heteroaryl and R<sup>6</sup> is hydrogen; R<sup>5</sup> and R<sup>6</sup> are independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl and aryl; or R<sup>5</sup> and R<sup>6</sup>, together with the nitrogen to which they are attached, form a ring comprising 4 to 6 carbon atoms or comprising 3 to 5 carbon atoms and one hetero moiety selected from the group consisting of -O- and -NR<sup>9</sup>-,  
 20 wherein R<sup>9</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl or phenyl; in the presence of a palladium catalyst and carbon monoxide to obtain an amide of formula 2:



; or


- 25 (ii) with an alcohol of the formula R<sup>10</sup>OH, wherein R<sup>10</sup> is C<sub>1</sub>-C<sub>6</sub> lower alkyl or C<sub>3</sub>-C<sub>6</sub> cycloalkyl, in the presence of a palladium catalyst and carbon monoxide to obtain the ester of formula 2A

-14-




followed by reacting the compound of 2A with an amine of formula  $\text{NHR}^5\text{R}^6$  to obtain the amide of formula 2;

(b) reacting the amide of formula 2 with a compound of formula 3




5

wherein  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  are as defined above and  $\text{R}^7$  is Cl or Br, in the presence of a strong base to obtain a compound of formula 4




10 (c)(i) converting a compound of formula 4 to a cyano compound of formula 5a




; or

(c)(ii) converting a compound of formula 4 or a cyano compound of formula 5a to an aldehyde of formula 5b




15 (d) reacting compound 5a or 5b with a piperidine derivative of formula 6





-15-

wherein L is a leaving group selected from the group consisting of Cl and Br, to obtain a ketone of formula 7a or an alcohol of formula 7b, respectively:




- 5            (e)(i) cyclizing a compound of formula 7a to obtain a compound of formula I wherein the dotted line represents a double bond; or
- (e)(ii) cyclizing a compound of formula 7b to obtain a compound of formula I wherein the dotted line represents a single bond.
  
- 10        2.        A process of claim 1 wherein R<sup>5</sup> is t-butylamine and R<sup>6</sup> is H, or R<sup>5</sup> and R<sup>6</sup> together are -(CH<sub>2</sub>)<sub>4</sub>.
  
3.        A process of claim 1 wherein the palladium catalyst is Pd(OAc)<sub>2</sub>/P(R<sup>10</sup>)<sub>3</sub>; (PPh<sub>3</sub>)<sub>2</sub>PdCl<sub>2</sub>; Pd(PPh<sub>3</sub>)<sub>4</sub>; (R<sup>10</sup>)<sub>3</sub>P/Pd<sub>2</sub>(dba)<sub>3</sub>; or
- 15        Pd/C, wherein Ac is acetyl, R<sup>10</sup> is C<sub>1</sub> to C<sub>6</sub> alkyl or aryl, Ph is phenyl, and dba is dibenzylidene acetone.
  
4.        A process of claim 1 wherein R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of chloro and bromo, and R<sup>3</sup> and R<sup>4</sup>
- 20        are each hydrogen.
  
5.        A process of claim 1 wherein R<sup>2</sup> and R<sup>4</sup> are independently selected from the group consisting of chloro and bromo, and R<sup>1</sup> and R<sup>3</sup>
- are each hydrogen.
  
- 25        6.        A process of claim 4 wherein R<sup>5</sup> is t-butylamine, R<sup>6</sup> is H, and the palladium catalyst is Pd(OAc)<sub>2</sub>/P(R<sup>11</sup>)<sub>3</sub> wherein Ac is acetyl and R<sup>11</sup> is C<sub>1</sub> to C<sub>6</sub> alkyl or aryl.


-16-

7. A process of claim 5 wherein  $R^5$  is t-butylamine,  $R^6$  is H, and the palladium catalyst is  $Pd(OAc)_2/P(R^{11})_3$  wherein Ac is acetyl and  $R^{11}$  is  $C_1$  to  $C_6$  alkyl or aryl.

- 5 8. A process of claim 1 comprising  
(a) reacting a compound of formula 1




- (i) with an amine of the formula  $NHR^5R^6$ , wherein  $R^5$  and  $R^6$  are as defined in claim 1, in the presence of a palladium catalyst and carbon monoxide to obtain an amide of formula 2:




; or

- (ii) with an alcohol of the formula  $R^{10}OH$ , wherein  $R^{10}$  is  $C_1$ - $C_6$  lower alkyl or  $C_3$ - $C_6$  cycloalkyl, in the presence of a palladium catalyst and carbon monoxide to obtain the ester of formula 2A




- 15 followed by reacting the compound of 2A with an amine of formula  $NHR^5R^6$  to obtain the amide of formula 2;

- (b) reacting the amide of formula 2 with a compound of formula 3




- 20 wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^7$  are as defined in claim 1, in the presence of a strong base to obtain a compound of formula 4




-17-

(c) converting a compound of formula 4 to a cyano compound of formula 5a




; or

(d) reacting compound 5a with a compound of formula 6



5

as defined in claim 1, to obtain a ketone of formula 7a:




(e)(i) cyclizing a compound of formula 7a to obtain a compound of formula I wherein the dotted line represents a double bond.

10


9. A process of claim 1 comprising:

(a) reacting a compound of formula 1



(i) with an amine of the formula  $\text{NHR}^5\text{R}^6$ , wherein  $\text{R}^5$  and  $\text{R}^6$  are as defined in claim 1, in the presence of a palladium catalyst and carbon monoxide to obtain an amide of formula 2:

15




; or

(ii) with an alcohol of the formula  $\text{R}^{10}\text{OH}$ , wherein  $\text{R}^{10}$  is  $\text{C}_1\text{-C}_6$  lower alkyl or  $\text{C}_3\text{-C}_6$  cycloalkyl, in the presence of a palladium catalyst and carbon monoxide to obtain the ester of formula 2A


20

-18-




followed by reacting the compound of 2A with an amine of formula  $\text{NHR}^5\text{R}^6$  to obtain the amide of formula 2;

(b) reacting the amide of formula 2 with a compound of formula 3




5

wherein  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$  and  $\text{R}^7$  are as defined in claim 1, in the presence of a strong base to obtain a compound of formula 4




10 (c)(i) converting a compound of formula 4 to a cyano compound of formula 5a




; or

(c)(ii) converting a compound of formula 4 or a cyano compound of formula 5a to an aldehyde of formula 5b




15

(d) reacting compound 5b with a compound of formula 6



-19-

as defined in claim 1, to obtain an alcohol of formula 7b:



(e)(ii) cyclizing a compound of formula 7b to obtain a compound of formula I wherein the dotted line represents a single bond.

## INTERNATIONAL SEARCH REPORT

National Application No

PCT/US 98/11403

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D401/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 151 423 A (J. J. PIWINSKI ET AL.) 29 September 1992 cited in the application see column 24, line 5 - column 26, line 42 see column 48, line 24 - column 49, line 36 ---	1,8,9
A	WO 95 10516 A (SCHERING CORP.) 20 April 1995 see page 49, line 6 - page 51, line 7 see page 61, line 11 - page 64, line 6 ---	1,8,9
A	WO 95 10515 A (SCHERING CORP.) 20 April 1995 see page 41, line 6 - page 43, line 7 see page 53, line 6 - page 55, line 23 --- -/--	1,8,9

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

27 August 1998

Date of mailing of the international search report

03/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/11403

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5151423 A	29-09-1992	US 5089496 A AT 114650 T AU 643946 B AU 5642090 A CA 2053903 A DE 69014393 D DE 69014393 T DK 396083 T EP 0396083 A EP 0471750 A ES 2064520 T IE 66392 B IL 94258 A JP 4504724 T KR 9509859 B NO 179674 B OA 9521 A WO 9013548 A US 5438062 A	18-02-1992 15-12-1994 02-12-1993 29-11-1990 02-11-1990 12-01-1995 01-06-1995 06-02-1995 07-11-1990 26-02-1992 01-02-1995 27-12-1995 07-10-1994 20-08-1992 29-08-1995 19-08-1996 15-11-1992 15-11-1990 01-08-1995
WO 9510516 A	20-04-1995	AU 7970394 A CA 2174104 A EP 0723540 A HU 76056 A JP 8510760 T US 5719148 A US 5714609 A US 5696121 A ZA 9407971 A	04-05-1995 20-04-1995 31-07-1996 30-06-1997 12-11-1996 17-02-1998 03-02-1998 09-12-1997 12-07-1996
WO 9510515 A	20-04-1995	AU 7930994 A CA 2174105 A EP 0723538 A HU 76066 A JP 8510759 T US 5721236 A US 5728703 A ZA 9407970 A	04-05-1995 20-04-1995 31-07-1996 30-06-1997 12-11-1996 24-02-1998 17-03-1998 12-07-1996
WO 9723478 A	03-07-1997	AU 1331197 A	17-07-1997

# INTERNATIONAL SEARCH REPORT

II  
of Application No  
PCT/US 98/11403

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	WO 97 23478 A (SCHERING CORP.) 3 July 1997 cited in the application see abstract -----	1